



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Global research priorities to better understand the burden of iatrogenic harm in primary care

Citation for published version:

World Health Organization's (WHO) Safer Primary Care Expert Working Group, Cresswell, KM, Panesar, SS, Salvilla, SA, Carson-Stevens, A, Larizgoitia, I, Donaldson, LJ, Bates, D & Sheikh, A 2013, 'Global research priorities to better understand the burden of iatrogenic harm in primary care: an international delphi exercise', *PLoS Medicine*, vol. 10, no. 11, pp. e1001554. <https://doi.org/10.1371/journal.pmed.1001554>

Digital Object Identifier (DOI):

[10.1371/journal.pmed.1001554](https://doi.org/10.1371/journal.pmed.1001554)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

PLoS Medicine

Publisher Rights Statement:

Copyright: © 2013 Cresswell et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Guidelines and Guidance

Global Research Priorities to Better Understand the Burden of Iatrogenic Harm in Primary Care: An International Delphi Exercise

Kathrin M. Cresswell¹, Sukhmeet S. Panesar², Sarah A. Salvilla², Andrew Carson-Stevens^{2,3}, Itziar Larizgoitia⁴, Liam J. Donaldson⁵, David Bates^{6,7,8}, Aziz Sheikh^{2,6,7*}, on behalf of the World Health Organization's (WHO) Safer Primary Care Expert Working Group[†]

1 The School of Health in Social Science, The University of Edinburgh, Edinburgh, United Kingdom, **2** Centre for Population Health Sciences, The University of Edinburgh, Edinburgh, United Kingdom, **3** Cochrane Institute of Primary Care & Public Health, School of Medicine, Cardiff University, Cardiff, United Kingdom, **4** WHO, Geneva, Switzerland, **5** Department of Surgery and Cancer, Imperial College London, London, United Kingdom, **6** Division of General Internal Medicine, Brigham and Women's Hospital, Boston, Massachusetts, United States of America, **7** Harvard Medical School, Boston, Massachusetts, United States of America, **8** Department of Health Policy and Management, Boston, Massachusetts, United States of America

Tackling the Provision of Unsafe Primary Care Internationally

It is now well established that medical errors are common and that these can result in considerable morbidity and mortality [1–3]. Much of this evidence, however, comes from hospital settings in industrialised countries where considerable progress has been made in describing the epidemiology of errors, understanding underlying contributing factors, and, more recently, taking steps to intervene to enhance patient safety [4].

In contrast, much less is known about the frequency of patient safety incidents and preventability of harm in primary care, particularly in low- and middle-income countries (Box 1). This is of concern as, in many parts of the world, primary care-based services now provide the first point of contact with health systems and often play a key role in coordinating more specialist care provision [5–8]. The increasing move to primary care-based health systems internationally [9,10] adds further impetus to the urgent need for research into the frequency and preventability of patient safety incidents, but this is complicated by the considerable variation in population needs, economic and political circumstances, structures of health systems, and manifestations of primary care globally.

In an attempt to support the development of a more comprehensive evidence-base, the World Health Organization (WHO) convened an international group of experts to discuss, debate, and advise on directions to bridge knowledge gaps around safe primary care, which would also serve to catalyse research in these areas internationally. A key strand of this foundational work was to identify a shared vision on relevant contexts of primary care and areas that would need further study to better understand the burden of harm in primary care settings internationally.

Developing Agreement on Primary Care Contexts and Priority Areas

We conducted a three-stage modified Delphi exercise, aiming to seek agreement on the most important contexts of primary care and the potential causes of patient safety incidents in different economic settings [11,12]. This exercise was undertaken during a two-day expert meeting in February 2012 at the WHO

headquarters in Geneva, Switzerland. AS, DB, and IL jointly chaired this meeting, which consisted of presentations, discussion groups, and plenary sessions focusing on understanding the challenges of assessing patient safety in primary care in low-, middle-, and high-income settings [13–15].

The Delphi technique has been widely used to help promote agreement amongst international experts. Key strengths of this process include the fact that it does not force consensus, but rather it can help to identify where agreement does and does not exist [16–19]. Although the general purpose and procedures have been retained in modified versions, some important differences in methods relate to: (1) approaches to managing interactions between experts, (2) the design of the initial item generation, and (3) feedback of individual scores [20–22].

Identifying Experts

We identified experts from academic, policy, and clinical backgrounds with expertise relating to patient safety in primary

Citation: Cresswell KM, Panesar SS, Salvilla SA, Carson-Stevens A, Larizgoitia I, et al. (2013) Global Research Priorities to Better Understand the Burden of Iatrogenic Harm in Primary Care: An International Delphi Exercise. *PLoS Med* 10(11): e1001554. doi:10.1371/journal.pmed.1001554

Published: November 19, 2013

Copyright: © 2013 Cresswell et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The meeting at which the data were collected was hosted by the Patient Safety Programme of WHO, Geneva, Switzerland. Limited resources were offered in some cases to cover travel expenses. Most participants or their affiliated institutions supported their own travel and attendance, with the University of Edinburgh, the Linnaeus Group Corporation, the Gulf Council Cooperation of States, and the PharmAccess Foundation sponsoring travel of various other participants. The University of Edinburgh was commissioned by WHO to support managing the meeting and providing the evidence base for the deliberations. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. AS is supported by The Commonwealth Fund, a private independent foundation based in New York City. The views presented here are those of the authors and not necessarily those of The Commonwealth Fund, its directors, officers, or staff.

Competing Interests: AS is a member of the Editorial Board of *PLOS Medicine*. All other authors have declared that no competing interests exist.

* E-mail: aziz.sheikh@ed.ac.uk

[†] Membership of the World Health Organization's (WHO) Safer Primary Care Expert Working Group is included in the Acknowledgments.

Provenance: Not commissioned; externally peer reviewed.

The Guidelines and Guidance section contains advice on conducting and reporting medical research.

Summary Points

- There is a need to identify and reach agreement on key foci for patient safety research in primary care contexts and understand how these priorities differ between low-, middle-, and high-income settings.
- We conducted a modified Delphi exercise, which was distributed to an international panel of experts in patient safety and primary care.
- Family practice and pharmacy were considered the main contexts on which to focus attention in order to advance patient safety in primary care across all income categories. Other clinical contexts prioritised included community midwifery and nursing in low-income countries and care homes in high-income countries.
- The sources of patient safety incidents requiring further study across all economic settings that were identified were communication between health care professionals and with patients, teamwork within the health care team, laboratory and diagnostic imaging investigations, issues relating to data management, transitions between different care settings, and chart/patient record completeness.
- This work lays the foundation for a range of research initiatives that aim to promote a more comprehensive appreciation of the burden of unsafe primary care, develop understanding of the main areas of risk, and identify interventions that can enhance the safety of primary care provision internationally.

care settings. This initial long-list was generated by drawing on existing WHO contacts, profiling academic institutions with relevant expertise in the area, identifying authors of published studies, and by searching Google Scholar. Selected experts were invited to participate at the meeting. Provisions were made to facilitate representation from low- and middle-income countries in different geographical regions, and by meeting travel costs where financial considerations were a possible barrier to attendance.

Generating Candidate Statements and Prioritisation Exercise

Participants were provided with a review of the literature surrounding the frequency of patient safety incidents, burden of harm, and preventability of these incidents in primary care. Data were collected in three iterative stages with corresponding data collection forms (see Text S1). The forms were distributed and collected face-to-face by members of the research team (KMC, SAS, SSP, and ACS). Opportunity for free text comments was provided throughout and each participant was assigned a number for anonymisation purposes.

The forms included a list of candidate areas identified from the literature, which were grouped into three sections with corresponding statements for low-, middle-, and high-income countries (see Text S1). After piloting, the list was shared with the experts at the beginning of day 1 of the meeting, asking participants to add additional items. This list was amended on the basis of participant feedback.

The amended list formed the basis for the second round of data collection. Here, experts were asked to score items in terms of importance (frequency of occurrence, severity of outcome, preventability, inequity of occurrence) on a 9-point Likert-type scale for each income category ranging from 1 = “not important”

Box 1. Glossary of Key Definitions

Primary care “Primary care is the provision of integrated, accessible health care services by clinicians who are accountable for:

addressing a large majority of personal health care needs developing a sustained partnership with patients practicing in the context of family and community” [10].

Patient safety “Patient safety is the reduction of risk of unnecessary harm associated with health care to an acceptable minimum. An acceptable minimum refers to the collective notions of given current knowledge, resources available and the context in which care was delivered weighed against the risk of non-treatment or other treatment” [28].

Harm “Harm implies impairment of structure or function of the body and/or any deleterious effect arising there from, including disease, injury, suffering, disability and death, and may be physical, social or psychological. Disease is a physiological or psychological dysfunction. Injury is damage to tissues caused by an agent or event and suffering is the experience of anything subjectively unpleasant. Suffering includes pain, malaise, nausea, depression, agitation, alarm, fear and grief. Disability implies any type of impairment of body structure or function, activity limitation and/or restriction of participation in society, associated with past or present harm” [28].

Low-, middle-, and high-income countries “For operational and analytical purposes, the World Bank’s main criterion for classifying economies is gross national income (GNI) per capita. In previous editions of our publications, this term was referred to as gross national product (GNP). Based on its GNI per capita, every economy is classified as low income, middle income (subdivided into lower middle and upper middle), or high income” [29].

to 9 = “extremely important” [15]. Data were independently scrutinised and transcribed by two members of the research team (KMC and ACS) into Microsoft Excel spread sheets and the median score for all items and the percentage agreement for items scoring 7, 8, or 9 (“usually important,” “very important,” and “extremely important,” respectively, i.e., the highest scores) were calculated.

The medians and percentage agreements obtained for each item were then included in the revised questionnaire that formed the basis for round 3 of data collection, giving participants the opportunity to revise their scoring on the basis of other participants’ rankings. The third questionnaire was distributed and collected at the end of day 1, followed by calculation of the percentage agreement with individual items.

Items with an agreement of >80% in each section at the end of the Delphi exercise were fed back to participants on day 2 of the meeting. Rather than feeding back the actual distribution of panellists’ ratings on the prior round, because of time and resource constraints the process fed back the median rating and percentage agreement of the prior round for the 7, 8, or 9 category as a proxy for the full distribution. This was followed by a plenary discussion, which gave participants the opportunity to collectively discuss emerging conclusions and recommendations. It enabled an




Figure 1. Flow-diagram of the three-stage modified Delphi process.
doi:10.1371/journal.pmed.1001554.g001

exploration of areas of convergence and divergence across countries and perspectives, giving participants the opportunity to air concerns and discuss potential next steps.

Overarching Considerations to Improve Understanding of the Extent of Unsafe Primary Care

We distributed 40 questionnaires in round 1. Of these, 37 questionnaires were completed. Reasons for non-completion were participants either failing to return the questionnaire ($n=1$) or leaving the meeting early ($n=2$). No potential participant explicitly refused to participate in the study. Of the 37 questionnaires distributed in round 2, 34 were returned (non-responses were again mainly because some participants needed to leave the meeting early ($n=3$)). In round 3, we distributed 30 questionnaires to the remaining participants (the other four had left the meeting),

all of which were completed. The overall response rate was therefore 30/40 (75%; see Figure 1). Key characteristics of those who completed all three rounds of the Delphi exercise are detailed in Table 1.

Overall, there was over 80% agreement across 15 items in low-income country contexts, 16 items in middle-income country contexts, and 16 items in high-income country contexts. Family practice and pharmacy were important primary care contexts across all income categories (Table 2). Additional contexts identified as warranting particular attention were community midwifery and nursing in low-income countries, and care homes in high-income countries (Table 2).

Tables 3 and 4 summarise the factors responsible for patient safety incidents that were identified as particularly needing further investigation across income settings. Important additional items in low- and middle-income settings included counterfeit drugs and errors in the execution of clinical tasks, whilst additional items in high-income settings were systems management and technology-related issues (Table 5).

Participants also prioritised the importance of cross-cutting systems' issues (Table 6). As can be seen, a range of interventional, regulatory, and methodological issues emerged; it is noteworthy that improved education and training for primary care workers received unanimous support.

Overall, we identified family practice and pharmacy as the main contexts to focus attention on in order to advance patient safety in primary care across all income categories. The sources of patient safety incidents requiring further study identified across all economic settings were communication between health care professionals and with patients, teamwork within the health care team, laboratory and diagnostic imaging investigations, issues relating to data management, transitions between different care settings, and chart/patient record completeness.

Strengths and Limitations of the Approach Employed

This work provides a foundation from which to focus efforts on how to better quantify the extent of iatrogenic harm in primary care and, in due course, to develop interventions to enhance the safety of primary care provision globally [5,6]. The exercise allowed us to identify areas for research into safer primary care, focusing on areas with the greatest propensity for harm and where

Table 1. Key participant characteristics of experts in all three rounds of the modified Delphi exercise.

Round	Gender	Professional Background	Number of Countries Represented	Income Settings Represented
1	8 female 29 male	8 academic (mostly doctors, some with pharmacy and nursing backgrounds) 7 non-for-profit research 10 health policy 7 clinical (all doctors) 5 academic/clinical	18	10 high-income 5 middle-income 3 low-income
2	8 female 26 male	8 academic 4 non-for-profit research 10 health policy 7 clinical 5 academic/clinical	17	9 high-income 5 middle-income 3 low-income
3	7 female 23 male	6 academic 3 non-for-profit research 10 health policy 6 clinical 5 academic/clinical	15	7 high-income 5 middle-income 3 low-income

doi:10.1371/journal.pmed.1001554.t001

Table 2. Primary care contexts that were considered to be important by over 80% of participants after round 3.

Primary care contexts prioritised across income settings
Family practice ^a
Pharmacy
Primary care contexts prioritised in low-income settings
Community midwifery
Community nursing
Primary care contexts prioritised in middle-income settings
Community nursing
Primary care contexts prioritised in high-income settings
Care homes

^aFamily practice was assumed to include general practice, outpatient paediatrics, and outpatient internal medicine.

doi:10.1371/journal.pmed.1001554.t002

prevention was considered feasible. Incorporating this exercise within a two-day face-to-face meeting helped to ensure good response rates retaining the majority of participants, and allowed feeding back results to participants in real time, taking their comments into account in analysis activities and questionnaire design. The iterative element of the exercise helped to inform the meeting's proceedings and ensuing discussions. We provided participants with the opportunity to discuss emerging areas of agreement, thereby contributing to the face validity of the results. The different backgrounds and expertise of participants created the opportunity to explore the specific challenges associated with primary care provision across a range of geographical and political contexts.

Our work also has some important limitations. The total number of participants was limited by resource constraints. Furthermore, as the focus was on policy and research deliberations, we had an over-representation of high-income country-based scholars and doctors, potentially influencing the results in favour of medical concerns important to those working in industrialised countries. That said, it is important to note that community pharmacy, nursing, and midwifery emerged as consistent priority areas. Some participants acknowledged that they had limited insights into provision of care in low-income settings, which may have influenced their ability to make informed judgements. Expansion of this exercise, involving additional experts from wider professional domains and world settings may therefore generate additional important insights.

Table 4. Five key causes of patient safety incidents that required further study across countries with different levels of income.

Problems resulting from poor communication and teamwork
Ordering and interpretation of diagnostic imaging and laboratory investigations
Issues relating to data management
Managing transitions between different levels of care
Completeness of patient records

doi:10.1371/journal.pmed.1001554.t004

Because of resource and time constraints in collating data over a narrow time window, we were unable to include a reminder of participants' own prior ratings, which may to an extent have been mitigated by the fact that all three rounds were conducted on the same day. We were also unable to provide additional information on distributions of ratings that could point to potentially diverging ratings and hence disagreements amongst experts that might not be reflected in the summary scores. We did however examine the raw data from each round for ambiguous items, but did not detect any such instances.

Finally, it should be noted that some known important patient safety issues in primary care such as injection safety were not included in this exercise because of a lack of specific expertise among participants, though it was noted as an important gap [23].

Implications and Unresolved Issues

Some of the existing knowledge underlying the measurement, causal factors, and interventions to enhance patient safety in primary care may be applicable across a wide array of income settings. This knowledge may be particularly relevant in relation to common contexts of care provision, including the central positioning of family practice and community pharmacy in health systems globally, although there may be other contextual and institutional factors at play that need to be better understood. The expert consultation strongly advocated the need for further research surrounding the frequency and preventability of patient safety incidents in primary care.

The discipline of patient safety is built on the premise that harm arises from errors in a multifactorial chain of events [24,25]. The underlying assumption is that if systems (i.e., organisations and networks of organisations) and working conditions within these

Table 3. Causes of patient safety incidents and their associated harm that were considered in need of further study by over 80% of participants after round 3 across all income settings.

Chart/patient record completeness
Communication between health care professionals in the same team
Communication between health care professionals from different teams
Data management
Laboratory investigations
Teamwork
Transitions between different levels of care
Wrong or missed diagnoses
Wrong treatment decision

doi:10.1371/journal.pmed.1001554.t003

Table 5. Items relating to causes of patient safety incidents and associated harm that were considered to be important by over 80% of participants after round 3 in low-, middle-, and high-income settings.

Causes of patient safety incidents in primary care prioritised in low-income settings
Counterfeit drugs
Execution of a clinical task (errors when performing clinical tasks due to lack of knowledge and/or skills)
Causes of patient safety incidents in primary care prioritised in middle-income settings
Communication between health care professionals and patients
Counterfeit drugs
Execution of a clinical task (errors when performing clinical tasks due to lack of knowledge and/or skills)
Higher-level systems management, e.g., human resources
Information technology and tools, e.g., checklists
Causes of patient safety incidents in primary care prioritised in high-income settings
Communication between health care professionals and patients
Diagnostic imaging
Higher-level systems management, e.g., human resources
Information technology and tools, e.g., checklists

doi:10.1371/journal.pmed.1001554.t005

organisations can be optimised, then the occurrence of adverse events is less likely. This systems approach is increasingly being applied as the patient safety culture of institutions and systems matures. Our findings support this trend, as reflected in the cross-cutting areas identified that all relate to improving ways of working collaboratively [26,27].

The work accomplished in this meeting can now be used as a starting point to inform and focus efforts in relation to epidemiological investigations that are urgently needed, particularly in low- and middle-income country contexts. These insights can then be used to develop interventions that aim to reduce risks of iatrogenic harm and improve health outcomes. Once tested, effective interventions need to be incorporated into local and international policy making in order to ensure that findings are effectively translated into practice.

Conclusions

Family practice and pharmacy were identified as important contexts across all income categories. Particular areas identified as warranting further investigation included communication between health care professionals and with patients, teamwork within the health care team, laboratory and diagnostic imaging investigations, issues relating to data management, transitions between different care settings, and chart/patient record completeness. The WHO will be issuing a roadmap within the next 12 months to ensure that the momentum from this important initiative is maintained.

Supporting Information

Text S1 Sample data collection form.
(DOCX)

Acknowledgments

We thank those who supported this work in various forms and are not included as authors. These include the members of the World Health

Table 6. Cross-cutting items that were considered to be important to focus on by over 80% of participants after round 3.

Education and training
Data collection methods
Developing policy to promote patient safety
Raising the public profile of patient safety
Greater clarity on definitions of errors in primary care
Facilitating learning from errors
Regulations to ensure that systems to improve patient safety are put into practice
Improved typologies/taxonomies (better ways of classifying errors in primary care)

doi:10.1371/journal.pmed.1001554.t006

Organization's (WHO) Safer Primary Care Expert Working Group: Carlos Aibar (Universidad de Zaragoza, Spain), Hamad Al-Bulushi (Ministry of Health, Oman), Buthaina Al-Mudaf (Ministry of Health, Kuwait), Hisham Aljadhey (King Saud University, Saudi Arabia), Fawzi Amin (Ministry of Health, Bahrain), Anthony Avery (The University of Nottingham, United Kingdom), Pierre Barker (Institute for Healthcare Improvement, United States of America), Jean Brami (Department of Evaluation, ANAES, France), Perpetual Chikobvu (University of the Free State, South Africa), Aneez Esmail (The University of Manchester, United Kingdom), John Hickner (Cleveland Clinic, United States of America), Neil Houston (Primary Care at Dollar Health Centre, United Kingdom), Tawfik Khoja (Health Ministers Council for Gulf Cooperation, Saudi Arabia), Maaike Langelaan (Netherlands Institute for Healthcare Research, Netherlands), Mondher Letaief (University Hospital of Monastir, Tunisia), Chaojie Liu (La Trobe University, Australia), Rajan Madhok (Global Association of Physicians of Indian Origin, United Kingdom/India), Meredith Makeham (University of New South Wales, Australia), Philippe Michel (Bordeaux University Hospital, France), Yakoub Neyaz (Ministry of Health, Saudi Arabia), Ludovic Reveiz Herault (AMRO/HSS/PR, United States of America), Gurdev Singh (Patient Safety Research Center, United States of America), Ranjit Singh (Patient Safety Research Center, United States of America), Andreas Soennichsen (Paracelsus Medical University, Austria), Nicole Spieker (PharmAccess Foundation, Netherlands), Hans Trier (Danish Society of Patient Safety, Denmark), Amardeep Thind (The University of Western Ontario, Canada), Nana Twum-Danso (Africa Operations at Institute for Healthcare Improvement, Ghana), Wim Verstappen (Radboud University Nijmegen Medical Centre, Netherlands), Katharine Wallis (University of Otago, New Zealand), Stuart Whittaker (COHSASA, South Africa), Benedetta Allegranzi (WHO), Edward Kelley (WHO), Angela Diane Lashofer (WHO), Shamsuzzoha Babar Syed (WHO), Antonio Villafaina (WHO), Yonatan Yohannes (WHO), Kevin Wang (WHO), Maria-Carmen Audera-Lopez (WHO), Marie-Paule Kieny (WHO). We are also most grateful to the institutions that sponsored participants to the meeting.

Author Contributions

Analyzed the data: KMC SSP SAS AC-S. Wrote the first draft of the manuscript: KMC AS. Contributed to the writing of the manuscript: SSP SAS AC-S IL IJD DB WHOSPCEWG. ICMJE criteria for authorship read and met: KMC SSP SAS AC-S IL IJD DB AS. Agree with manuscript results and conclusions: KMC SSP SAS AC-S IL IJD DB AS WHOSPCEWG. KMC and SSP led the drafting of the protocol, which received critical input from SAS and AC-S.

References

1. The Health Foundation (2011) Improving safety in primary care. Available: <http://www.health.org.uk/publications/improving-safety-in-primary-care-research-scan/>. Accessed 7 September 2013.
2. Fiss T, Dreier A, Meinke C, van den Berg N, Ritter CA, et al. (2011) Frequency of inappropriate drugs in primary care: analysis of a sample of immobile patients who received periodic home visits. *Age Ageing* 40: 66–73.
3. Woolf SH, Kuzel AJ, Dovey SM, Phillips RL, Jr. (2004) A string of mistakes: the importance of cascade analysis in describing, counting, and preventing medical errors. *Ann Fam Med* 2: 317–26.
4. Makeham M, Dovey S, Runciman W, Larizgiotia I (2008) Methods and measures used in primary care patient safety research. Available: http://www.who.int/patientsafety/research/methods_measures/makeham_dovey_full.pdf. Accessed 7 September 2013.
5. Donaldson SL (2005) World alliance for patient safety: forward programme. Geneva: WHO.
6. Bates DW, Larizgiotia I, Prasopa-Plaizier N, Jha AK (2009) Global priorities for patient safety research. *BMJ* 338: b1775.
7. Starfield B (1998) Primary care: balancing health needs, services and technology. New York: Oxford University Press.
8. Reerink IH, Sauerborn R (1996) Quality of primary health care in developing countries: recent experiences and future directions. *Int J Qual Health Care* 8: 131–9.
9. WHO (1978) Declaration of Alma-Ata. In International Conference on Primary Health Care. Alma-Ata, USSR: WHO.
10. Institute of Medicine (1996) Primary care: America's health in a new era. Available: http://www.nap.edu/openbook.php?record_id=5152&page=27. Accessed 7 September 2013.
11. Mir G, Salway S, Kai J, Karlens S, Bhopal R, et al. (2013) Principles for research on ethnicity and health: the Leeds Consensus Statement. *Eur J Public Health* 23: 504–510.
12. Avery AJ, Savelyich BS, Sheikh A, Cantrill J, Morris CJ, et al. (2005) Identifying and establishing consensus on the most important safety features of GP computer systems: e-Delphi study. *Inform Prim Care* 13: 3–12.
13. Jones J, Hunter D (1995) Consensus methods for medical and health services research. *BMJ* 311: 376–80.
14. Cuhls K (2012) Delphi method. Available: www.unido.org/fileadmin/import/16959_DelphiMethod.pdf. Accessed 7 September 2013.
15. Brook RH, Chassin MR, Fink A, Solomon DH, Koseoff J, et al. (1986) A method for the detailed assessment of the appropriateness of medical technologies. *Int J Technol Assess Health Care* 2: 53–63.
16. Elwyn G, O'Connor A, Stacey D, Volk R, Edwards A, et al. (2006) Developing a quality criteria framework for patient decision aids: online international Delphi consensus process. *BMJ* 333: 417–23.
17. Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, et al. (2005) Global prevalence of dementia: a Delphi consensus study. *Lancet* 366: 2112–7.
18. Shekelle P, Woolf SH, Eccles M, Grimshaw J (1999) Clinical guidelines: developing guidelines. *BMJ* 318: 593–6.
19. Shekelle P (2004) The appropriateness method. *Med Decis Making* 24: 228–31.
20. Stewart J, O'Halloran C, Harrigan P, Spencer JA, Barton JR, et al. (1999) Identifying appropriate tasks for the preregistration year: modified Delphi technique. *BMJ* 319: 224–9.
21. Mullen PM (2003) Delphi: myths and reality. *J Health Organ Manag* 17: 37–52.
22. De Meyrick J (2012) The Delphi method and health research. *Health Education* 103: 7–16.
23. WHO (2012) Injection safety. Available: http://www.who.int/injection_safety/en/. Accessed 7 September 2013.
24. Institute of Medicine (1999) To err is human: building a safer health system. Washington (D.C.): National Academy Press.
25. Reason J (2000) Human error: models and management. *BMJ* 320: 768–70.
26. Preston C, Cheater F, Baker R (1999) Left in limbo: patients' views on care processes across the primary/secondary interface. *Quality in Health Care* 8: 16–21.
27. Roland M (1995) R&D priorities in relation to the interface between primary and secondary care. Report to the Central Research and Development Committee. Leicester: Department of Health.
28. WHO (2009) More than words. Conceptual Framework for the International Classification for Patient Safety. Available: http://www.who.int/patientsafety/implementation/taxonomy/icps_technical_report_en.pdf. Accessed 7 September 2013.
29. World Bank (2012) How we classify countries. Available: <http://data.worldbank.org/about/country-classifications>. Accessed 7 September 2013.